

Exclusive Semileptonic Rare Decays $B \rightarrow (K, K^*)\ell^+\ell^-$ in Supersymmetric Theories

Qi-Shu YAN*

Physics Department of Tsinghua University, P.R.China

Chao-Shang HUANG[†], LIAO Wei[‡], Shou-Hua ZHU[§]

Institute of Theoretical Physics,
the Chinese Academy Science, P.R.China

Abstract

The invariant mass spectrum, forward-backward asymmetry, and lepton polarizations of the exclusive processes $B \rightarrow K(K^*)\ell^+\ell^-$, $\ell = \mu, \tau$ are analyzed under supersymmetric context. Special attention is paid to the effects of neutral Higgs bosons (NHBs). Our analysis shows that the branching ratio of the process $B \rightarrow K\mu^+\mu^-$ can be quite largely modified by the effects of neutral Higgs bosons and the forward-backward asymmetry would not vanish. For the process $B \rightarrow K^*\mu^+\mu^-$, the lepton transverse polarization is quite sensitive to the effects of NHBs, while the invariant mass spectrum, forward-backward asymmetry, and lepton longitudinal polarization are not. For both $B \rightarrow K\tau^+\tau^-$ and $B \rightarrow K^*\tau^+\tau^-$, the effects of NHBs are quite significant. The partial decay widths of these processes are also analyzed, and our analysis manifest that even taking into account the theoretical uncertainties in calculating weak form factors, the effects of NHBs could make SUSY shown up.

PACS numbers: 13.20.He, 13.25.Hw, 12.60.Jv

* E-mail : yanqs@itp.ac.cn; qsyang@mail.tsinghua.edu.cn

[†] E-mail : csh@itp.ac.cn

[‡] E-mail: liaow@itp.ac.cn

[§] E-mail: huald@itp.ac.cn

1 Introduction

The inclusive rare processes $b \rightarrow X_s \ell^+ \ell^-$, $\ell = e, \mu, \tau$ have been intensively studied in literatures [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. As one of flavor changing neutral current processes, it is sensitive to fine structure of the standard model and to the possible new physics as well, and is expected to shed light on the existence of new physics before the possible new particles are produced at colliders.

It is well known that invariant mass spectrum, forward-backward asymmetries (FBAs), and lepton polarizations are important observables to probe new physics, while the first two observables are mostly analyzed. About lepton polarizations, it is known that due to the smallness of the mass of it, therefore electron polarizations are very difficult to be measured experimentally. So only the lepton polarizations of muon and tau are considered in literatures [10, 12, 13, 14]. The longitudinal polarization of tau in $B \rightarrow X_s \tau^+ \tau^-$ has been calculated in standard model (SM) and several new physics scenarios [10]. For $B \rightarrow X_s l^+ l^-$ ($l = \mu, \tau$), the polarizations of lepton in SM are analyzed in [12] and it is pointed out that for the μ channel, the only significant component is the longitudinal polarization (P_L), while all three components are sizable in the τ channel. The analysis has been extended to supersymmetric models (SUSY) and a CP softly broken two Higgs doublet model in refs. [13] and [14] respectively. The reference [5] also gives a general model-independent analysis of the lepton polarization asymmetries in the process $B \rightarrow X_s \tau^+ \tau^-$ and it is found that the contribution from $C_{LRLR} + C_{LRRL}$ is much larger than other scalar-type interactions.

Compared with the inclusive processes $B \rightarrow X_s \ell^+ \ell^-$, $\ell = e, \mu, \tau$, the theoretical study of the exclusive processes $B \rightarrow K(K^*) \ell^+ \ell^-$ is relatively hard. For inclusive semileptonic decays of B, the decay rates can be calculated in heavy quark effective theory (HQET) [15]. However, for exclusive semileptonic decays of B, to make theoretical predictions, additional knowledge of decay form factors is needed, which is related with the calculation of hadronic transition matrix elements. Hadronic transition matrix elements depend on the non-perturbative properties of QCD, and can only be reliably calculated by using a nonperturbative method. The form factors for B decay into $K^{(*)}$ have been computed with different methods such as quark models [16], SVZ QCD sum rules [17], light cone sum rules (LCSRs) [18, 19, 20, 21, 22]. Compared to the lattice approach which mainly deal with the form factors at small recoil, the QCD sum rules on the light-cone can complementarily provide information of the form factors at smaller values of $\hat{s}$. And they are consistent with perturbative QCD and the heavy quark limit. In this work, we will use the weak decay form factors calculated by using the technique of the light cone QCD sum rules and given in [23].

A upper limit on the branching ratio of $B^0 \rightarrow K^{0*} \mu^+ \mu^-$ has been recently given by CLEO [24]:

$$BR(B^0 \rightarrow K^{0*} \mu^+ \mu^-) < 4.0 \times 10^{-6}, \quad (1.1)$$

and they will be precisely measured at B factories, these exclusive processes are quite worthy of intensive study and have attracted many attentions [23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36]. In reference [23], by using improved theoretical calculations of the decay form factors in the light cone QCD sum rule approach, dilepton invariant mass spectra and the FBAs of these exclusive decays are analyzed in the standard model and a number of popular variants of the supersymmetric models. However, as the author claimed, the effects of neutral Higgs exchanges are neglected. For exclusive processes, as pointed out in [36], the polarization

asymmetries of μ and τ for $B \rightarrow K^* \mu^+ \mu^-$ and $B \rightarrow K^* \tau^+ \tau^-$ are also accessible at the B-factories under construction. In reference [34], the lepton polarizations and CP violating effects in $B \rightarrow K^* \tau^+ \tau^-$ are analyzed in SM and two Higgs doublet models.

As pointed in refs. [3, 4], in two-Higgs-doublet models and SUSY models, neutral Higgs boson could contribute largely to the inclusive processes $B \rightarrow X_s \ell^+ \ell^-$, $\ell = \mu, \tau$ and greatly modify the branching ratio and FBA in the large $\tan\beta$ case. The effects of neutral Higgs in the 2HDM to polarizations of τ in $B \rightarrow K \tau^+ \tau^-$ are analyzed in [35], and it was found that polarizations of the charged final lepton are very sensitive to the $\tan\beta$.

In this paper, we will investigate the exclusive decay $B \rightarrow K(K^*) \ell^+ \ell^-$, $\ell = \mu, \tau$ in SUSY models. We shall evaluate branching ratios and FBAs with emphasis on the effects of neutral Higgs and analyze lepton polarizations in MSSM. According to the analysis of [29], different sources of the vector current could manifest themselves in different regions of phase space, for the very low $\hat{s}$ the photonic penguin dominates, while the Z penguin and W box becomes important towards high $\hat{s}$. In order to search the regions of $\hat{s}$ where neutral Higgs bosons could contribute large, we analyze the partial decay widths of these two processes. Beside that they are accessible to B factories, our motivation also bases on the fact that to the inclusive processes $B \rightarrow X_s \ell^+ \ell^-$, $\ell = \mu, \tau$, neutral Higgs could make quite a large contributions at certain large $\tan\beta$ regions of parameter space in SUSY models, since part of supersymmetric contributions is proportional to $\tan^3\beta$ [4]. Such regions considerably exist in SUGRA and M-theory inspired models [37]. We also analyze the effects of neutral Higgs to the position of the zero value of the FBA. Our results show that the branching ratio of the process $B \rightarrow K \mu^+ \mu^-$ can be quite largely modified by the effects of neutral Higgs bosons and the FBA would not vanish. Because the FBA for $B \rightarrow K \ell^+ \ell^-$ ($\ell = \mu, \tau$) vanishes if the contributions of NHBs vanish. The contributions of NHBs can be large enough to be observed only in SUSY and/or 2HDM with large $\tan\beta$, and a non-zero FBA for $B \rightarrow K \ell^+ \ell^-$ would signal the existence of new physics. For the process $B \rightarrow K^* \mu^+ \mu^-$, the lepton transverse polarization is quite sensitive to the effects of NHBs, while the invariant mass spectrum, FBA, and lepton longitudinal polarization are not. For both $B \rightarrow K \tau^+ \tau^-$ and $B \rightarrow K^* \tau^+ \tau^-$, the effects of NHBs are quite significant. Our analysis manifest that even taking into account the theoretical uncertainties in calculating weak form factors, the effects of NHBs could show SUSY up. In brief, our analysis manifest that effects of NHBs is quite remarkable in some regions of parameter space of SUSY, even for the process $B \rightarrow K \mu^+ \mu^-$.

The paper is organized as follows. In the section 2, the effective Hamiltonian is presented and the form factors given by using light cone sum rule method are briefly discussed. Basic formula of observables are introduced in section 3. Section 4 is devoted to the numerical analysis. In section 5 we make discussions and conclusions.

2 Effective Hamiltonian and Form Factors

By integrating out the degrees of heavy freedom from the full theory, MSSM, at electroweak(EW) scale, we can get the effective Hamiltonian describing the rare semileptonic decay $b \rightarrow s \ell^+ \ell^-$:

$$\mathcal{H}_{\text{eff}} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \left(\sum_{i=1}^{10} C_i(\mu) O_i(\mu) + \sum_{i=1}^{10} C_{Q_i}(\mu) Q_i(\mu) \right), \quad (2.1)$$

where the first ten operators and Wilson coefficients at EW scale can be found in [8, 38]¹, and last ten operators and Wilson coefficients which represent the contributions of neutral Higgs can be found in [4].

With the renormalization group equations to resum the QCD corrections, Wilson coefficients at energy scale $\mu = m_b$ are evaluated. Theoretical uncertainties related to renormalization-scale can be substantially reduced when the next-leading-logarithm corrections are included [39].

The above Hamiltonian leads to the following free quark decay amplitude:

$$\begin{aligned} \mathcal{M}(b \rightarrow s \ell^+ \ell^-) = & -\frac{G_F \alpha}{\sqrt{2} \pi} V_{ts}^* V_{tb} \left\{ C_9^{\text{eff}} [\bar{s} \gamma_\mu L b] [\bar{\ell} \gamma^\mu \ell] + C_{10} [\bar{s} \gamma_\mu L b] [\bar{\ell} \gamma^\mu \gamma_5 \ell] \right. \\ & \left. - 2 \hat{m}_b C_7^{\text{eff}} \left[\bar{s} i \sigma_{\mu\nu} \frac{\hat{q}^\nu}{\hat{s}} R b \right] [\bar{\ell} \gamma^\mu \ell] + C_{Q1} [\bar{s} R b] [\bar{\ell} \ell] + C_{Q2} [\bar{s} R b] [\bar{\ell} \gamma_5 \ell] \right\} \end{aligned} \quad (2.2)$$

where C_9^{eff} is defined as [40, 41]

$$C_9^{\text{eff}}(\mu, \hat{s}) = C_9(\mu) + Y(\mu, \hat{s}) + \frac{3\pi}{\alpha^2} C(\mu) \sum_{V_i=\psi(1s), \dots, \psi(6s)} \kappa_i \frac{\Gamma(V_i \rightarrow \ell^+ \ell^-) m_{V_i}}{m_{V_i}^2 - \hat{s} m_B^2 - i m_{V_i} \Gamma_{V_i}} \quad (2.3)$$

where $\hat{s} = s/m_b^2$, $s=q^2$, $C(\mu) = (3C_1 + C_2 + 3C_3 + C_4 + 3C_5 + C_6)$, and

$$\begin{aligned} Y(\mu, \hat{s}) = & g(\hat{m}_c, \hat{s}) C(\mu) \\ & - \frac{1}{2} g(1, \hat{s}) (4C_3 + 4C_4 + 3C_5 + C_6) - \frac{1}{2} g(0, \hat{s}) (C_3 + 3C_4) \\ & - \frac{2}{9} (3C_3 + C_4 + 3C_5 + C_6) . \end{aligned} \quad (2.4)$$

where the function $g(\hat{m}_c, \hat{s})$ comes from one loop contributions of four-quark operators and is defined by

$$g(z, \hat{s}) = -\frac{4}{9} \ln z^2 + \frac{8}{27} + \frac{16}{9} \frac{z^2}{\hat{s}} - \begin{cases} \frac{2}{9} \sqrt{1 - \frac{4z^2}{\hat{s}}} \left(2 + \frac{4z^2}{\hat{s}} \right) \left[\ln \left(\frac{1 + \sqrt{1 - 4z^2/\hat{s}}}{1 - \sqrt{1 - 4z^2/\hat{s}}} \right) - i\pi \right], & 4z^2 < \hat{s} \\ \frac{4}{9} \sqrt{\frac{4z^2}{\hat{s}} - 1} \left(2 + \frac{4z^2}{\hat{s}} \right) \arctan \left(\frac{1}{\sqrt{4z^2/\hat{s} - 1}} \right), & 4z^2 > \hat{s} \end{cases} \quad (2.5)$$

The last terms in (2.3) are nonperturbative effects from $(\bar{c}c)$ resonance contributions, while the phenomenological factors κ_i can be fixed from the processes [23] $B \rightarrow K^{(*)} V_i \rightarrow K^{(*)} \ell^+ \ell^-$ and as given in the Table. 1.

Exclusive decays $B \rightarrow (K, K^*) \ell^+ \ell^-$ are described in terms of matrix elements of the quark operators in Eq. (2.2) over meson states, which can be parametrized in terms of form factors.

¹In our previous papers, e.g., [3, 4], we follow the convention of ref. [1] for the indices of operators as well as Wilson coefficients. In this paper, we use more popular conventions (see, e.g., [39]). That is, $O_8 \rightarrow O_9$ and $O_9 \rightarrow O_{10}$.

For the process $B \rightarrow K\ell^+\ell^-$, the non-vanishing matrix elements are ($q = p_B - p$)

$$\langle K(p)|\bar{s}\gamma_\mu b|B(p_B)\rangle = f_+(s) \left\{ (p_B + p)_\mu - \frac{m_B^2 - m_K^2}{s} q_\mu \right\} + \frac{m_B^2 - m_K^2}{s} f_0(s) q_\mu, \quad (2.6)$$

and

$$\begin{aligned} \langle K(p)|\bar{s}\sigma_{\mu\nu}q^\nu(1 + \gamma_5)b|B(p_B)\rangle &= \langle K(p)|\bar{s}\sigma_{\mu\nu}q^\nu b|B(p_B)\rangle \\ &= i \left\{ (p_B + p)_\mu s - q_\mu(m_B^2 - m_K^2) \right\} \frac{f_T(s)}{m_B + m_K}. \end{aligned} \quad (2.7)$$

While for $B \rightarrow K^*\ell^+\ell^-$, related transition matrix elements are

$$\begin{aligned} \langle K^*(p)|(V - A)_\mu|B(p_B)\rangle &= -i\epsilon_\mu^*(m_B + m_{K^*})A_1(s) + i(p_B + p)_\mu(\epsilon^*p_B) \frac{A_2(s)}{m_B + m_{K^*}} \\ &\quad + i q_\mu(\epsilon^*p_B) \frac{2m_{K^*}}{s} (A_3(s) - A_0(s)) + \epsilon_{\mu\nu\rho\sigma}\epsilon^{*\nu}p_B^\rho p^\sigma \frac{2V(s)}{m_B + m_{K^*}}. \end{aligned} \quad (2.8)$$

and

$$\begin{aligned} \langle K^*|\bar{s}\sigma_{\mu\nu}q^\nu(1 + \gamma_5)b|B(p_B)\rangle &= i\epsilon_{\mu\nu\rho\sigma}\epsilon^{*\nu}p_B^\rho p^\sigma 2T_1(s) \\ &\quad + T_2(s) \left\{ \epsilon_\mu^*(m_B^2 - m_{K^*}^2) - (\epsilon^*p_B)(p_B + p)_\mu \right\} \\ &\quad + T_3(s)(\epsilon^*p_B) \left\{ q_\mu - \frac{s}{m_B^2 - m_{K^*}^2} (p_B + p)_\mu \right\} \end{aligned} \quad (2.9)$$

Where ϵ_μ is polarization vector of the vector meson K^* . By means of the equation of motion, one obtains several relations between form factors

$$\begin{aligned} A_3(s) &= \frac{m_B + m_{K^*}}{2m_{K^*}} A_1(s) - \frac{m_B - m_{K^*}}{2m_{K^*}} A_2(s), \\ A_0(0) &= A_3(0), \\ \langle K^*|\partial_\mu A^\mu|B\rangle &= 2m_{K^*}(\epsilon^*p_B)A_0(s), \\ T_1(0) &= T_2(0). \end{aligned} \quad (2.10)$$

κ	J/Ψ	Ψ'
K	2.70	3.51
K^*	1.65	2.36

Table 1: *Fudge factors in $B \rightarrow K^{(*)}J/\Psi, \Psi' \rightarrow K^{(*)}\ell^+\ell^-$ decays calculated using the LCSR form factors.*

All signs are defined in such a way as to render the form factors real and positive. The physical range in $\hat{s}$ extends from $\hat{s}_{\min} = 4\hat{m}_\ell^2$ to $\hat{s}_{\max} = (1 - \hat{m}_{K,K^*})^2$.

The calculation of the form factors given above is a real task, and one has to rely on certain approximate methods. We use the results calculated by using technique of LCSRs and given in [23]. And form factors can be parametrized as

$$F(\hat{s}) = F(0) \exp(c_1 \hat{s} + c_2 \hat{s}^2 + c_3 \hat{s}^3). \quad (2.11)$$

The parameterization formula works within 1% accuracy for $s < 15 \text{ GeV}^2$ and can avoid the spurious singularities at $s = m_B^2$. Related parameters is given in the Table. 4 of [23]

3 Formula of Observables

In this section we provide formula of experimental observables, which include dilepton invariant mass spectrum, FBA, and lepton polarizations.

From eqs. (2.2 - 2.8), it is straightforward to obtain the matrix element of $B \rightarrow K(K^*)l^+l^-$ as follows.

$$\mathcal{M} = -\frac{G_F \alpha}{2\sqrt{2}\pi} V_{ts}^* V_{tb} m_B \left[\mathcal{T}_\mu^1 (\bar{\ell} \gamma^\mu \ell) + \mathcal{T}_\mu^2 (\bar{\ell} \gamma^\mu \gamma_5 \ell) + \mathcal{S} (\bar{\ell} \ell) \right], \quad (3.1)$$

where for $B \rightarrow K \ell^+ \ell^-$,

$$\mathcal{T}_\mu^1 = A'(\hat{s}) \hat{p}_\mu, \quad (3.2)$$

$$\mathcal{T}_\mu^2 = C'(\hat{s}) \hat{p}_\mu + D'(\hat{s}) \hat{q}_\mu, \quad (3.3)$$

$$\mathcal{S} = \mathcal{S}_1(\hat{s}) \quad (3.4)$$

and for $B \rightarrow K^* \ell^+ \ell^-$,

$$\mathcal{T}_\mu^1 = A(\hat{s}) \epsilon_{\mu\rho\alpha\beta} \epsilon^{*\rho} \hat{p}_B^\alpha \hat{p}_{K^*}^\beta - iB(\hat{s}) \epsilon_\mu^* + iC(\hat{s}) (\epsilon^* \cdot \hat{p}_B) \hat{p}_\mu, \quad (3.5)$$

$$\mathcal{T}_\mu^2 = E(\hat{s}) \epsilon_{\mu\rho\alpha\beta} \epsilon^{*\rho} \hat{p}_B^\alpha \hat{p}_{K^*}^\beta - iF(\hat{s}) \epsilon_\mu^* + iG(\hat{s}) (\epsilon^* \cdot \hat{p}_B) \hat{p}_\mu + iH(\hat{s}) (\epsilon^* \cdot \hat{p}_B) \hat{q}_\mu, \quad (3.6)$$

$$\mathcal{S} = i2\hat{m}_{K^*} (\epsilon^* \cdot \hat{p}_B) \mathcal{S}_2(\hat{s}) \quad (3.7)$$

with $p \equiv p_B + p_{K,K^*}$. Note that, using the equation of motion for lepton fields, the terms in $\hat{q}_\mu$ in $\mathcal{T}_\mu^1$ vanish.

The auxiliary functions above are defined as

$$A'(\hat{s}) = C_9^{\text{eff}}(\hat{s}) f_+(\hat{s}) + \frac{2\hat{m}_b}{1 + \hat{m}_K} C_7^{\text{eff}} f_T(\hat{s}), \quad (3.8)$$

$$C'(\hat{s}) = C_{10} f_+(\hat{s}), \quad (3.9)$$

$$D'(\hat{s}) = C_{10} f_-(\hat{s}) - \frac{1 - \hat{m}_K^2}{2\hat{m}_\ell(\hat{m}_b - \hat{m}_s)} C_{Q2} f_0(\hat{s}), \quad (3.10)$$

$$\mathcal{S}_1(\hat{s}) = \frac{1 - \hat{m}_K^2}{(\hat{m}_b - \hat{m}_s)} C_{Q1} f_0(\hat{s}) , \quad (3.11)$$

$$A(\hat{s}) = \frac{2}{1 + \hat{m}_{K^*}} C_9^{\text{eff}}(\hat{s}) V(\hat{s}) + \frac{4\hat{m}_b}{\hat{s}} C_7^{\text{eff}} T_1(\hat{s}) , \quad (3.12)$$

$$B(\hat{s}) = (1 + \hat{m}_{K^*}) \left[C_9^{\text{eff}}(\hat{s}) A_1(\hat{s}) + \frac{2\hat{m}_b}{\hat{s}} (1 - \hat{m}_{K^*}) C_7^{\text{eff}} T_2(\hat{s}) \right] , \quad (3.13)$$

$$C(\hat{s}) = \frac{1}{1 - \hat{m}_{K^*}^2} \left[(1 - \hat{m}_{K^*}) C_9^{\text{eff}}(\hat{s}) A_2(\hat{s}) + 2\hat{m}_b C_7^{\text{eff}} \left(T_3(\hat{s}) + \frac{1 - \hat{m}_{K^*}^2}{\hat{s}} T_2(\hat{s}) \right) \right] , \quad (3.14)$$

$$E(\hat{s}) = \frac{2}{1 + \hat{m}_{K^*}} C_{10} V(\hat{s}) , \quad (3.15)$$

$$F(\hat{s}) = (1 + \hat{m}_{K^*}) C_{10} A_1(\hat{s}) , \quad (3.16)$$

$$G(\hat{s}) = \frac{1}{1 + \hat{m}_{K^*}} C_{10} A_2(\hat{s}) , \quad (3.17)$$

$$\begin{aligned} H(\hat{s}) &= \frac{C_{10}}{\hat{s}} [(1 + \hat{m}_{K^*}) A_1(\hat{s}) - (1 - \hat{m}_{K^*}) A_2(\hat{s}) - 2\hat{m}_{K^*} A_0(\hat{s})] \\ &\quad + \frac{\hat{m}_{K^*}}{\hat{m}_\ell(\hat{m}_b + \hat{m}_s)} A_0(\hat{s}) C_{Q2} , \end{aligned} \quad (3.18)$$

$$\mathcal{S}_2(\hat{s}) = -\frac{1}{(\hat{m}_b + \hat{m}_s)} A_0(\hat{s}) C_{Q1} . \quad (3.19)$$

where

$$f_0(\hat{s}) = \frac{1}{1 - \hat{m}_K^2} [\hat{s} f_-(\hat{s}) + (1 - \hat{m}_K^2) f_+(\hat{s})] \quad (3.20)$$

and to get the auxiliary functions given above, we have used equations of motion

$$q^\mu (\bar{\psi}_1 \gamma_\mu \psi_2) = (m_1 - m_2) \bar{\psi}_1 \psi_2 , \quad (3.21)$$

$$q^\mu (\bar{\psi}_1 \gamma_\mu \gamma_5 \psi_2) = -(m_1 + m_2) \bar{\psi}_1 \gamma_5 \psi_2 . \quad (3.22)$$

The contributions of NHBs have been incorporated in the terms of $\mathcal{S}_1(\hat{s})$, $D'(\hat{s})$, $H(\hat{s})$ and $\mathcal{S}_2(\hat{s})$. It is remarkable that the contributions of NHBs in $D'(\hat{s})$ and $H(\hat{s})$ are proportional to the inverse mass of the lepton, and for the case $l = \mu$, the effects of NHBs can be manifested through these terms.

A phenomenological effective Hamiltonian is recently given in [30]. If ignoring tensor type interactions in the phenomenological Hamiltonian (it is shown that physical observables are not sensitive to the presence of tensor type interactions [6]), it is easy to verify that the matrix element of $B \rightarrow K^{(*)} \ell^+ \ell^-$ can always be expressed as the form of the equation (3.1) with the

auxiliary functions defined as

$$A'(\hat{s}) = w_{c_1} f_+(\hat{s}) - \frac{w_{c_9} + w_{c_{10}}}{1 + \hat{m}_K} f_T(\hat{s}) , \quad (3.23)$$

$$C'(\hat{s}) = w_{c_2} f_+(\hat{s}) , \quad (3.24)$$

$$D'(\hat{s}) = w_{c_2} f_-(\hat{s}) - \frac{1 - \hat{m}_K^2}{2\hat{m}_\ell(\hat{m}_b - \hat{m}_s)} w_{c_6} f_0(\hat{s}) , \quad (3.25)$$

$$\mathcal{S}_1(\hat{s}) = \frac{1 - \hat{m}_K^2}{(\hat{m}_b - \hat{m}_s)} w_{c_5} f_0(\hat{s}) , \quad (3.26)$$

$$A(\hat{s}) = \frac{2}{1 + \hat{m}_{K^*}} w_{c_1} V(\hat{s}) - \frac{2}{\hat{s}} (w_{c_9} + w_{c_{10}}) T_1(\hat{s}) , \quad (3.27)$$

$$B(\hat{s}) = -(1 + \hat{m}_{K^*}) \left[w_{c_3} A_1(\hat{s}) + \frac{1}{\hat{s}} (1 - \hat{m}_{K^*}) (w_{c_9} + w_{c_{10}}) T_2(\hat{s}) \right] , \quad (3.28)$$

$$C(\hat{s}) = -\frac{1}{1 - \hat{m}_{K^*}^2} \left[(1 - \hat{m}_{K^*}) w_{c_3}(\hat{s}) A_2(\hat{s}) \right. \\ \left. + (w_{c_9} - w_{c_{10}}) \left((1 + \hat{m}_{K^*}) T_3(\hat{s}) + \frac{1 - \hat{m}_{K^*}^2}{\hat{s}} T_2(\hat{s}) \right) \right] , \quad (3.29)$$

$$E(\hat{s}) = \frac{2}{1 + \hat{m}_{K^*}} w_{c_2} V(\hat{s}) , \quad (3.30)$$

$$F(\hat{s}) = -(1 + \hat{m}_{K^*}) w_{c_4} A_1(\hat{s}) , \quad (3.31)$$

$$G(\hat{s}) = -\frac{1}{1 + \hat{m}_{K^*}} w_{c_4} A_2(\hat{s}) , \quad (3.32)$$

$$H(\hat{s}) = -\frac{2\hat{m}_{K^*}}{\hat{s}} w_{c_4} (A_3(\hat{s}) - A_0(\hat{s})) + \frac{\hat{m}_{K^*}}{\hat{m}_\ell(\hat{m}_b + \hat{m}_s)} w_{c_8} A_0(\hat{s}) , \quad (3.33)$$

$$\mathcal{S}_2(\hat{s}) = -\frac{1}{(\hat{m}_b + \hat{m}_s)} w_{c_7} A_0(\hat{s}) , \quad (3.34)$$

where

$$w_{c_1} = \frac{1}{4} (C_{LL} + C_{LR} + C_{RL} + C_{RR}) , \quad (3.35)$$

$$w_{c_2} = \frac{1}{4} (-C_{LL} + C_{LR} - C_{RL} + C_{RR}) , \quad (3.36)$$

$$w_{c_3} = \frac{1}{4} (-C_{LL} - C_{LR} + C_{RL} + C_{RR}) , \quad (3.37)$$

$$w_{c_4} = \frac{1}{4}(C_{LL} - C_{LR} - C_{RL} + C_{RR}) , \quad (3.38)$$

$$w_{c_5} = \frac{1}{4}(C_{LRLR} + C_{RLLR} + C_{LRRL} + C_{RLRL}) , \quad (3.39)$$

$$w_{c_6} = \frac{1}{4}(C_{LRLR} + C_{RLLR} - C_{LRRL} - C_{RLRL}) , \quad (3.40)$$

$$w_{c_7} = \frac{1}{4}(C_{LRLR} - C_{RLLR} + C_{LRRL} - C_{RLRL}) , \quad (3.41)$$

$$w_{c_8} = \frac{1}{4}(C_{LRLR} - C_{RLLR} - C_{LRRL} + C_{RLRL}) , \quad (3.42)$$

$$w_{c_9} = m_b C_{BR} , \quad (3.43)$$

$$w_{c_{10}} = m_s C_{SL} . \quad (3.44)$$

In the above equations C_{LL}, C_{LR} etc are defined in ref. [6]. Therefore our formula given below can also be used to make model independent phenomenological analysis, if using Eqs. ((3.23)-(3.34)) instead of Eqs. ((3.8)-(3.19)).

Keeping the lepton mass, we find the double differential decay widths Γ^K and Γ^{K^*} for the decays $B \rightarrow K \ell^+ \ell^-$ and $B \rightarrow K^* \ell^+ \ell^-$, respectively, as

$$\begin{aligned} \frac{d^2 \Gamma^K}{d\hat{s} d\hat{u}} &= \frac{G_F^2 \alpha^2 m_B^5}{2^{11} \pi^5} |V_{ts}^* V_{tb}|^2 \\ &\times \left\{ (|A'|^2 + |C'|^2)(\lambda - \hat{u}^2) + |\mathcal{S}_1|^2(\hat{s} - 4\hat{m}_\ell^2) + \text{Re}(\mathcal{S}_1 A'^*) 4\hat{m}_\ell \hat{u} \right. \\ &\left. + |C'|^2 4\hat{m}_\ell^2 (2 + 2\hat{m}_K^2 - \hat{s}) + \text{Re}(C' D'^*) 8\hat{m}_\ell^2 (1 - \hat{m}_K^2) + |D'|^2 4\hat{m}_\ell^2 \hat{s} \right\} , \quad (3.45) \end{aligned}$$

$$\begin{aligned} \frac{d^2 \Gamma^{K^*}}{d\hat{s} d\hat{u}} &= \frac{G_F^2 \alpha^2 m_B^5}{2^{11} \pi^5} |V_{ts}^* V_{tb}|^2 \\ &\times \left\{ \frac{|A|^2}{4} (\hat{s}(\lambda + \hat{u}^2) + 4\hat{m}_\ell^2 \lambda) + \frac{|E|^2}{4} (\hat{s}(\lambda + \hat{u}^2) - 4\hat{m}_\ell^2 \lambda) + |\mathcal{S}_2|^2 (\hat{s} - 4\hat{m}_\ell^2) \lambda \right. \\ &\left. + \frac{1}{4\hat{m}_{K^*}^2} \left[|B|^2 (\lambda - \hat{u}^2 + 8\hat{m}_{K^*}^2 (\hat{s} + 2\hat{m}_\ell^2)) + |F|^2 (\lambda - \hat{u}^2 + 8\hat{m}_{K^*}^2 (\hat{s} - 4\hat{m}_\ell^2)) \right] \right. \\ &\left. - 2\hat{s}\hat{u} [\text{Re}(BE^*) + \text{Re}(AF^*)] + \frac{2\hat{m}_\ell \hat{u}}{\hat{m}_{K^*}} [\text{Re}(\mathcal{S}_2 B^*) (\hat{s} + \hat{m}_{K^*}^2 - 1) + \text{Re}(\mathcal{S}_2 C^*) \lambda] \right. \\ &\left. + \frac{\lambda}{4\hat{m}_{K^*}^2} \left[|C|^2 (\lambda - \hat{u}^2) + |G|^2 (\lambda - \hat{u}^2 + 4\hat{m}_\ell^2 (2 + 2\hat{m}_{K^*}^2 - \hat{s})) \right] \right. \\ &\left. - \frac{1}{2\hat{m}_{K^*}^2} [\text{Re}(BC^*) (1 - \hat{m}_{K^*}^2 - \hat{s}) (\lambda - \hat{u}^2) \right] \end{aligned}$$

$$\begin{aligned}
& +\text{Re}(FG^*) \left[(1 - \hat{m}_{K^*}^2 - \hat{s})(\lambda - \hat{u}^2) + 4\hat{m}_\ell^2 \lambda \right] \\
& - 2 \frac{\hat{m}_\ell^2}{\hat{m}_{K^*}^2} \lambda \left[\text{Re}(FH^*) - \text{Re}(GH^*)(1 - \hat{m}_{K^*}^2) \right] + |H|^2 \frac{\hat{m}_\ell^2}{\hat{m}_{K^*}^2} \hat{s} \lambda \Big\} .
\end{aligned} \tag{3.46}$$

Here the kinematic variables $(\hat{s}, \hat{u})$ are defined as

$$\hat{s} = \hat{q}^2 = (\hat{p}_+ + \hat{p}_-)^2 , \tag{3.47}$$

$$\hat{u} = (\hat{p}_B - \hat{p}_-)^2 - (\hat{p}_B - \hat{p}_+)^2 \tag{3.48}$$

which are bounded as

$$(2\hat{m}_\ell)^2 \leq \hat{s} \leq (1 - \hat{m}_{K,K^*})^2 , \tag{3.49}$$

$$-\hat{u}(\hat{s}) \leq \hat{u} \leq \hat{u}(\hat{s}) , \tag{3.50}$$

with $\hat{m}_\ell = m_\ell/m_B$ and

$$\hat{u}(\hat{s}) = \sqrt{\lambda(1 - 4\frac{\hat{m}_\ell^2}{\hat{s}})} , \tag{3.51}$$

$$\lambda = 1 + \hat{m}_{K,K^*}^4 + \hat{s}^2 - 2\hat{s} - 2\hat{m}_{K,K^*}^2(1 + \hat{s}) , \tag{3.52}$$

$$\mathcal{D} = \sqrt{1 - \frac{4\hat{m}_\ell^2}{\hat{s}}} . \tag{3.53}$$

Note that the variable $\hat{u}$ corresponds to θ , the angle between the momentum of the B -meson and the positively charged lepton ℓ^+ in the dilepton CMS frame, through the relation $\hat{u} = -\hat{u}(\hat{s}) \cos \theta$ [42].

Integrating over $\hat{u}$ in the kinematic region given in Eq. (3.50) we get the formula of dilepton invariant mass spectra (IMS)

$$\frac{d\Gamma^K}{d\hat{s}} = \frac{G_F^2 \alpha^2 m_B^5}{2^{10} \pi^5} |V_{ts}^* V_{tb}|^2 \hat{u}(\hat{s}) D^K \tag{3.54}$$

$$\begin{aligned}
D^K = & (|A'|^2 + |C'|^2) \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} \right) + |\mathcal{S}_1|^2 (\hat{s} - 4\hat{m}_\ell^2) \\
& + |C'|^2 4\hat{m}_\ell^2 (2 + 2\hat{m}_K^2 - \hat{s}) + \text{Re}(C' D'^*) 8\hat{m}_\ell^2 (1 - \hat{m}_K^2) + |D'|^2 4\hat{m}_\ell^2 \hat{s} ,
\end{aligned} \tag{3.55}$$

$$\frac{d\Gamma^{K^*}}{d\hat{s}} = \frac{G_F^2 \alpha^2 m_B^5}{2^{10} \pi^5} |V_{ts}^* V_{tb}|^2 \hat{u}(\hat{s}) D^{K^*} \tag{3.56}$$

$$\begin{aligned}
D^{K^*} = & \frac{|A|^2}{3} \hat{s} \lambda (1 + 2\frac{\hat{m}_\ell^2}{\hat{s}}) + |E|^2 \hat{s} \frac{\hat{u}(\hat{s})^2}{3} + |\mathcal{S}_2|^2 (\hat{s} - 4\hat{m}_\ell^2) \lambda \\
& + \frac{1}{4\hat{m}_{K^*}^2} \left[|B|^2 \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} + 8\hat{m}_{K^*}^2 (\hat{s} + 2\hat{m}_\ell^2) \right) + |F|^2 \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} + 8\hat{m}_{K^*}^2 (\hat{s} - 4\hat{m}_\ell^2) \right) \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{\lambda}{4\hat{m}_{K^*}^2} \left[|C|^2 \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} \right) + |G|^2 \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} + 4\hat{m}_\ell^2(2 + 2\hat{m}_{K^*}^2 - \hat{s}) \right) \right] \\
& - \frac{1}{2\hat{m}_{K^*}^2} \left[\text{Re}(BC^*) \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} \right) (1 - \hat{m}_{K^*}^2 - \hat{s}) \right. \\
& \left. + \text{Re}(FG^*) \left(\left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} \right) (1 - \hat{m}_{K^*}^2 - \hat{s}) + 4\hat{m}_\ell^2 \lambda \right) \right] \\
& - 2 \frac{\hat{m}_\ell^2}{\hat{m}_{K^*}^2} \lambda \left[\text{Re}(FH^*) - \text{Re}(GH^*) (1 - \hat{m}_{K^*}^2) \right] + \frac{\hat{m}_\ell^2}{\hat{m}_{K^*}^2} \hat{s} \lambda |H|^2. \tag{3.57}
\end{aligned}$$

Both distributions agree with the ones obtained in [23, 36], if $C_{Q1,2}$ are set to zero. The differential FBA is defined as

$$A_{FB}(s) = \frac{-\int_0^{u(\hat{s})} dz \frac{d\Gamma}{dsdu} + \int_{-u(\hat{s})}^0 du \frac{d\Gamma}{dsdu}}{\int_0^{u(\hat{s})} dz \frac{d\Gamma}{dsdu} + \int_{-u(\hat{s})}^0 du \frac{d\Gamma}{dsdu}}.$$

For $B \rightarrow K\ell^+\ell^-$ decays it reads as follows

$$\frac{d\mathcal{A}_{\text{FB}}^K}{d\hat{s}} D^K = -2\hat{m}_\ell \hat{u}(\hat{s}) \text{Re}(\mathcal{S}_1 A'^*) \tag{3.58}$$

For $B \rightarrow K^*\ell^+\ell^-$ decays it reads as follows

$$\begin{aligned}
\frac{d\mathcal{A}_{\text{FB}}^{K^*}}{d\hat{s}} D^{K^*} &= \hat{u}(\hat{s}) \left\{ \hat{s} [\text{Re}(BE^*) + \text{Re}(AF^*)] \right. \\
&\left. + \frac{\hat{m}_\ell}{\hat{m}_{K^*}} [\text{Re}(\mathcal{S}_2 B^*) (1 - \hat{s} - \hat{m}_{K^*}^2) - \text{Re}(\mathcal{S}_2 C^*) \lambda] \right\} \tag{3.59}
\end{aligned}$$

We can read from (3.58), the FBA of the process $B \rightarrow K\ell^+\ell^-$ does not vanish when the contributions of NHB are taken into account. With it, our analysis below also show that the contributions of NHBs can even be accessible in B factories.

The lepton polarization can be defined as follows

$$\frac{d\Gamma(\vec{n})}{ds} = \frac{1}{2} \left(\frac{d\Gamma}{ds} \right)_0 \left[1 + (P_L \vec{e}_L + P_N \vec{e}_N + P_T \vec{e}_T) \cdot \vec{n} \right], \tag{3.60}$$

where the subscript "0" corresponds to the unpolarized width, and P_L , P_T , and P_N , correspond to the longitudinal, transverse and normal components of the polarization vector, respectively.

For the process $B \rightarrow K\ell^-\ell^+$, the P_L^K , P_T^K , and P_N^K , are derived respectively as

$$P_L^K D^K = \frac{4}{3} \mathcal{D} \left\{ \lambda \text{Re}(A' C'^*) - 3\hat{m}_\ell (1 - \hat{m}_K^2) \text{Re}(C'^* \mathcal{S}_1) - 3\hat{m}_\ell \hat{s} \text{Re}(D'^* \mathcal{S}_1) \right\}, \tag{3.61}$$

$$P_N^K D^K = \frac{\pi\sqrt{\hat{s}}\hat{u}(\hat{s})}{2} \left\{ -\text{Im}(A'\mathcal{S}_1^*) + 2\hat{m}_\ell \text{Im}(C'D'^*) \right\}, \quad (3.62)$$

$$P_T^K D^K = \frac{-\pi\sqrt{\lambda}}{\sqrt{\hat{s}}} \left\{ \hat{m}_\ell \left[(1 - \hat{m}_{K^*}^2) \text{Re}(A'C'^*) \right. \right. \\ \left. \left. + \hat{s} \text{Re}(A'D'^*) \right] + \frac{(\hat{s} - 4\hat{m}_\ell^2)}{2} \text{Re}(C'\mathcal{S}_1^*) \right\}. \quad (3.63)$$

D^K is defined in Eq. (3.55). For the process $B \rightarrow K^* \ell^- \ell^+$, the $P_L^{K^*}$, $P_T^{K^*}$, and $P_N^{K^*}$, are derived respectively as

$$P_L^{K^*} D^{K^*} = \mathcal{D} \left\{ \frac{2\hat{s}\lambda}{3} \text{Re}(AE^*) + \frac{(\lambda + 12\hat{m}_{K^*}^2)}{3\hat{m}_{K^*}^2} \text{Re}(BF^*) \right. \\ \left. - \frac{\lambda(1 - \hat{m}_{K^*}^2 - \hat{s})}{3\hat{m}_{K^*}^2} \text{Re}(BG^* + CF^*) + \frac{\lambda^2}{3\hat{m}_{K^*}^2} \text{Re}(CG^*) \right. \\ \left. + \frac{2\hat{m}_\ell\lambda}{\hat{m}_{K^*}} \left[\text{Re}(F\mathcal{S}_2^*) - \hat{s} \text{Re}(H\mathcal{S}_2^*) - (1 - \hat{m}_{K^*}^2) \text{Re}(G\mathcal{S}_2^*) \right] \right\}, \quad (3.64)$$

$$P_N^{K^*} D^{K^*} = \frac{-\pi\sqrt{\hat{s}}\hat{u}(\hat{s})}{4\hat{m}_K} \left\{ \frac{\hat{m}_\ell}{\hat{m}_{K^*}} \left[\text{Im}(FG^*)(1 + 3\hat{m}_{K^*}^2 - s) \right. \right. \\ \left. \left. + \text{Im}(FH^*)(1 - \hat{m}_{K^*}^2 - s) - \text{Im}(GH^*)\lambda \right] \right. \\ \left. + 2\hat{m}_{K^*}\hat{m}_\ell [\text{Im}(BE^*) + \text{Im}(AF^*)] \right. \\ \left. - (1 - \hat{m}_{K^*}^2 - \hat{s}) \text{Im}(B\mathcal{S}_2^*) + \lambda \text{Im}(C\mathcal{S}_2^*) \right\}, \quad (3.65)$$

$$P_T^{K^*} D^{K^*} = \frac{\pi\sqrt{\lambda}\hat{m}_\ell}{4\sqrt{\hat{s}}} \left\{ 4\hat{s} \text{Re}(AB^*) \right. \\ \left. + \frac{(1 - \hat{m}_{K^*}^2 - \hat{s})}{\hat{m}_{K^*}^2} \left[-\text{Re}(BF^*) + (1 - \hat{m}_{K^*}^2) \text{Re}(BG^*) + \hat{s} \text{Re}(BH^*) \right] \right. \\ \left. + \frac{\lambda}{\hat{m}_{K^*}^2} \left[\text{Re}(CF^*) - (1 - \hat{m}_{K^*}^2) \text{Re}(CG^*) - \hat{s} \text{Re}(CH^*) \right] \right. \\ \left. + \frac{(\hat{s} - 4\hat{m}_\ell^2)}{\hat{m}_{K^*}\hat{m}_\ell} \left[(1 - \hat{m}_{K^*}^2 - \hat{s}) \text{Re}(F\mathcal{S}_2^*) - \lambda \text{Re}(G\mathcal{S}_2^*) \right] \right\}. \quad (3.66)$$

D^{K^*} is defined by Eq.(3.57).

m_b	4.8 GeV
m_c	1.4 GeV
m_s	0.2 GeV
m_μ	0.11 GeV
m_τ	1.78 GeV
M_B	5.28 GeV
M_K	0.49 GeV
M_{K^*}	0.89 GeV
$M_{J/\psi}(M_{\psi'})$	3.10(3.69) GeV
Γ_B	4.22×10^{-13} GeV
$\Gamma_{J/\psi}(\Gamma_{\psi'})$	$8.70(27.70) \times 10^{-5}$ GeV
$\Gamma(J/\psi \rightarrow \ell^+ \ell^-)$	5.26×10^{-6} GeV
$\Gamma(\psi' \rightarrow \ell^+ \ell^-)$	2.14×10^{-6} GeV
G_F	1.17×10^{-5} GeV ⁻²
α^{-1}	129
$ V_{ts}^* V_{tb} $	0.0385

Table 2: *Values of the input parameters used in our numerical analysis.*

C_1	C_2	C_3	C_4	C_5	C_6	C_7^{eff}	C_9	C_{10}	C
-0.248	+1.107	+0.011	-0.026	+0.007	-0.031	-0.313	+4.344	-4.669	+0.362

Table 3: *Wilson coefficients of the SM used in the numerical analysis.*

4 numerical analysis

Parameters used in our analysis are list in Table 2. Considering that the branching ratios of $B \rightarrow K \ell^+ \ell^-$ and $B \rightarrow K^* \ell^+ \ell^-$ are not very sensitive to the mass of m_b , we neglect the difference between the pole mass and running mass of b quark.

The Wilson coefficients in the SM used in the numerical analysis is given in the Table 3. C_7^{eff} is defined as

$$C_7^{\text{eff}} = C_7 - C_5/3 - C_6. \quad (4.67)$$

$C_{Q_{1,2}}$ come from exchanging NHBs and are proportional to $\tan^3 \beta$ in some regions of the parameter space in SUSY models. According to the analysis in [4, 37], the necessary conditions for the large contributions of NHBs include: (i) the ratio of vacuum expectation value, $\tan \beta$, should be large, (ii) the mass values of the lighter chargino and the lighter stop should not be too large (say less than 120 GeV), (iii) mass splitting of charginos and stops should be large, which also indicate large mixing between stop sector and chargino sector. As the conditions are satisfied, the process $B \rightarrow X_s \gamma$ will impose a constraint on C_7 . It is well known that this process puts a very stringent constraint on the possible new physics and that SUSY can

SUSY models	R_7	R_9	R_{10}	C_{Q_1}	C_{Q_2}
SUSY I	-1.2	1.1	0.8	0.0	0.0
SUSY II	-1.2	1.1	0.8	6.5(16.5)	-6.5(-16.5)
SUSY III	1.2	1.1	0.8	1.2(4.5)	-1.2(-4.5)

Table 4: *Wilson coefficients of the SUSY used in our numerical analysis. R_i means C_i/C_i^{SM} . SUSY I corresponds to the regions where SUSY can destructively contribute and can change the sign of C_7 , but the contributions of NHBs are neglected. SUSY II corresponds to the regions where $\tan\beta$ is large and the masses of superpartners are relatively small. SUSY III corresponds to the regions where $\tan\beta$ is large but the masses of superpartners are relatively large. In the last two cases the effects of NHBs are taken into account. The contributions of NHBs are settled to be different for both the case $\ell = \mu$ and $\ell = \tau$, since $C_{Q_{1,2}}$ are proportional to the mass of lepton. The values in bracket are for the case $\ell = \tau$.*


2

contribute destructively when the signature of the Higgs mass term μ is minus. There exist considerable regions of SUSY parameter space in which NHBs can largely contribute to the process $b \rightarrow s\ell^+\ell^-$ while the constraint of $b \rightarrow s\gamma$ is respected (i.e., the signature of the Wilson coefficient C_7 is changed from positive to negative). When the masses of SUSY particle are relatively heavy (say, 450 GeV), there are still significant regions in the parameter space of SUSY models in which NHBs could contribute largely. However, at these cases C_7 does not change its sign, because contributions of charged Higgs and charginos cancel with each other. We will see it is hopeful to distinguish these two kinds of regions of SUSY parameter space through observing $B \rightarrow K^{(*)}\ell^+\ell^-$.


As pointed out in [3, 4], the contribution of NHBs is proportional to the lepton mass, therefore for $\ell = e$, contributions of NHBs can be safely neglected. While for cases $\ell = \mu$ and $\ell = \tau$, the contributions of NHBs can be considerably large. To investigate the effects of NHBs in SUSY models, we take typical values of $C_{7,9,10}$ and $C_{Q_{1,2}}$ as given in Table 6. The SUSY model without considering the effects of NHBs (SUSY I in Table 4) is given as a reference frame so that could the effects of NHBs be shown in high relief.

Numerical results are shown in Figs. 1-4. In Fig. 1(a), the IMS of $B \rightarrow K\mu^+\mu^-$ is depicted. We see that at the high $\hat{s}$ regions, NHBs greatly modify the spectrum. While at the low $\hat{s}$ region, the effects of NHBs become weak. In Fig. 1(b), the FBA of the $B \rightarrow K\mu^+\mu^-$ is presented. Fig. 1(b) shows that the average FBA in $B \rightarrow K\mu^+\mu^-$ is 0.02. To measure an asymmetry A of a decay with the branching ratio Br at the $n\sigma$ level, the required number of events is $N = n^2/(BrA^2)$. For $B \rightarrow K\mu^+\mu^-$, the average FBA is 0.02 or so, the required number of events is 10^{12} or so. Therefore it is hard to observe the derivation of FBA from the SM. In Fig. 1(c) and Fig. 1(d), the longitudinal and transverse polarizations are given. The effect of NHBs to the longitudinal polarization is weak but the effect to the transverse is remarkable.


In Fig. 2(a) and Fig. 2(b) the IMS and FBA of $B \rightarrow K\tau^+\tau^-$ are presented respectively. For SUSY II, the effects of NHBs to IMS is quite manifest, and the average FBA can reach 0.1. For SUSY III, the average FBA can reach 0.3. Therefore, in order to observe FBA, the




(a)



(b)




(c)




(d)


Figure 1: The $IMS(a)$, $FBA(b)$, $P_L(c)$, and $P_T(d)$ of the process $B \rightarrow K\mu^+\mu^-$. The solid line, dashed line, dot line and dashed-dot line represent the SM, SUSY I, SUSY II, SUSY III respectively. Both the total (SD+LD) and the pure SD contributions are shown in order to compare.




(a)



(b)



(c)



(d)

Figure 2: The $IMS(a)$, $FBA(b)$, $P_L(c)$, and $P_T(d)$ of the process $B \rightarrow K\tau^+\tau^-$. The line conventions are the same as given in the legend of Fig 1.

required number of events should be 10^9 or so and 10^8 , respectively, so that in B factories, say LHCb, these two cases are accessible. In Fig. 2(c) and Fig. 2(d), the longitudinal and transverse polarizations are drawn respectively. The effects of NHBs are also very obvious.


Figs. 3 and 4 are devoted to the decay $B \rightarrow K^* l^+ l^-$. In Fig. 3, the IMS, FBA, and polarizations of $B \rightarrow K^* \mu^+ \mu^-$ are given. We see that this process is not as much as sensitive to the effect of NHB as $B \rightarrow K \mu^+ \mu^-$. However, the contribution of NHBs will increase the part with positive FBA and will be helpful to determine the zero point of FBA. Fig. 3(d) depicts the transverse polarization of the $B \rightarrow K^* \mu^+ \mu^-$, and the effect of NHBs is quite obvious. The zero point of the FBA can be slightly modified as shown in Figure 3(b) due to the contributions of NHBs.

In Fig. 4, the IMS, FBA, longitudinal and transverse polarizations of the $B \rightarrow K^* \tau^+ \tau^-$ are depicted. The effect of NHBs does show in great relief. It is worth to note that IMS, FBA, and lepton polarizations for $B \rightarrow K^* \ell^+ \ell^-$ in MSSM without including the contributions of NHBs are also significantly different from those in SM, while for $B \rightarrow K \ell^+ \ell^-$ they have little differences from those in SM. Therefore, compared to the process $B \rightarrow K \ell^+ \ell^-$, more precise measurements for $B \rightarrow K^* \ell^+ \ell^-$ are needed in order to single out the contributions of NHBs.


Normal polarizations for both $B \rightarrow K \ell^+ \ell^-$ and $B \rightarrow K^* \ell^+ \ell^-$ are small and can be neglected because the imaginary parts of Wilson coefficients are small in SUSY models without CP violating phases which are implicitly assumed in the paper.

The behavior of IMS(a), FBA(b), P_L (c), and P_T (d) shown Figs 1-4 can be understood with the formula given in the Section 3. With Eqs. (3.55), (3.10) and (3.11), we see that the contributions of NHBs are contained in the terms of $\mathcal{S}_1$ and D' . At the high $\hat{s}$ regions, it is these two terms which are important. This explained the behavior of IMS given in (a) of Fig. 1 and Fig. 2. The Eq. (3.58) shows that the FBA is proportional to the mass of the lepton. For the case $B \rightarrow K \mu^+ \mu^-$, due to smallness of the mass μ , the FBA does not vanish but is hard to be measured. While for the case $B \rightarrow K \tau^+ \tau^-$, the mass τ is quite large and observing FBA is relatively easy. For SUSY II, though the numerator of FBA is comparatively large, the large IMS suppresses the value of FBA; for SUSY III, the numerator is relatively small, but the FBA do demonstrate the effects of NHBs more manifestly, as shown in Fig. 2(b) due to smallness of IMS. The Eqs. (3.63) and (3.64) show that for the case $\ell = \mu$, the contributions of NHBs to P_N, P_T are suppressed by the mass of μ . But for the case $\ell = \tau$, the contributions of NHBs become quite manifest both for SUSY II and SUSY III. The term with D' in Eq. (3.63) will change its sign when there exists relatively not too small contributions of NHBs, the fact deduced from Eq. (3.10), that explains why the sign of P_T is changed. The difference between the case SUSY II and SUSY III is small, the reason is just the same as stated in the analysis of FBA.


Since the terms incorporating the contributions of NHBs is proportional to λ as shown in Eq. (3.57), which approaches zero at high $\hat{s}$ regions; while at small $\hat{s}$ regions, the effects of NHBs are dwarfed by the other contributions. Therefore, only when C_{Q_i} are quite large could effects of NHBs be manifest, as shown in Fig. 3(a) and Fig. 4(a). According to the Eq. (3.59), at high $\hat{s}$ regions, the effects of NHBs would be suppressed by λ and $1 - \hat{s} - \hat{m}_{K^*}^2$. The same suppression mechanism exists for P_L . This suppression mechanism explains the fact that the processes $B \rightarrow K^* \ell^+ \ell^-$ are not sensitive to the effects of NHBs. However, when there exist large contributions of NHBs, the sign of P_T will be changed, as indicated in both Fig. 3(d) and Fig. 4(d).




(a)



(b)




(c)




(d)


Figure 3: *The IMS(a), FBA(b), P_L (c), and P_T (d) of the process $B \rightarrow K^*\mu^+\mu^-$. The line conventions are the same as given in the legend of Fig 1.*




(a)



(b)



(c)



(d)

Figure 4: The $IMS(a)$, $FBA(b)$, $P_L(c)$, and $P_T(d)$ of the process $B \rightarrow K^* \tau^+ \tau^-$. The line conventions are the same as given in the legend of Fig 1.

model		A	B	C	D	E	tot(SD)	tot(SD+LD)
SM	LCSR	0.353	54.707	0.032	4.566	0.076	0.573	59.736
	SVZ	0.215	22.918	0.015	1.593	0.026	0.299	24.767
SUSY I	LCSR	0.425	54.723	0.037	4.576	0.086	0.675	59.847
	SVZ	0.179	22.910	0.011	1.586	0.019	0.236	24.704
SUSY II	LCSR	0.556	54.865	0.131	4.833	0.849	2.067	61.233
	SVZ	0.348	23.009	0.068	1.726	0.321	1.002	25.473
SUSY III	LCSR	0.429	54.727	0.040	4.584	0.109	0.717	59.889
	SVZ	0.181	22.912	0.012	1.590	0.028	0.255	24.723

Table 5: Partial decay widths for $B \rightarrow K\mu^+\mu^-$. LCSR means the approach light-cone QCD sum rules, SVZ means the SVZ QCD sum rule [17]. Character A means the region $(\hat{s}_0, (\hat{m}_\psi - \hat{\delta})^2)$, B $((\hat{m}_\psi - \hat{\delta})^2, (\hat{m}_\psi + \hat{\delta})^2)$, C $((\hat{m}_\psi + \hat{\delta})^2, (\hat{m}_{\psi'} - \hat{\delta})^2)$, D $((\hat{m}_{\psi'} - \hat{\delta})^2, (\hat{m}_{\psi'} + \hat{\delta})^2)$ and E $((\hat{m}_{\psi'} + \hat{\delta})^2, \hat{s}_{\max}^2)$. The unit is $\Gamma_B \times 10^{-6}$, which is 4.22×10^{-19} GeV. δ is selected to be 0.2 GeV. $\hat{\delta}$ is normalized with M_B

model		A	B	C	D	E	tot(SD)	tot(SD+LD)
SM	LCSR	0.930	83.257	0.141	9.976	0.258	1.882	94.562
	SVZ	2.943	111.278	0.147	7.504	0.137	3.639	122.008
SUSY I	LCSR	1.627	83.402	0.198	10.085	0.330	2.915	95.64
	SVZ	4.517	111.423	0.183	7.552	0.149	5.291	123.825
SUSY II	LCSR	1.178	83.431	0.234	10.164	0.352	2.677	95.360
	SVZ	2.801	111.292	0.156	7.525	0.145	3.522	121.918
SUSY III	LCSR	1.631	83.407	0.201	10.092	0.334	2.938	95.664
	SVZ	4.518	111.425	0.184	7.553	0.150	5.296	123.830

Table 6: Partial decay widths for $B \rightarrow K^*\mu^+\mu^-$. Other conventions can be found in Table 5.

model		A'	B'	tot(SD)	tot(SD+LD)
SM	LCSR	1.884	0.094	0.132	1.978
	SVZ	0.659	0.036	0.054	0.695
SUSY I	LCSR	1.884	0.086	0.131	1.970
	SVZ	0.655	0.025	0.038	0.680
SUSY II	LCSR	2.022	1.496	1.674	3.519
	SVZ	0.726	0.552	0.637	1.278
SUSY III	LCSR	1.874	0.094	0.129	1.968
	SVZ	0.651	0.026	0.035	0.677

Table 7: Partial decay widths of $B \rightarrow K\tau^+\tau^-$. Character A' means $(\hat{s}_0, (\hat{m}_\psi - \hat{\delta})^2)$, B' means $((\hat{m}_{\psi'} + \hat{\delta})^2, \hat{s}_{\max})$. The unit is $\Gamma_B \times 10^{-6}$, which is 4.22×10^{-19} GeV.

model		A'	B'	tot(SD)	tot(SD+LD)
SM	LCSR	4.045	0.096	0.183	4.141
	SVZ	3.029	0.048	0.102	3.076
SUSY I	LCSR	4.088	0.173	0.327	4.261
	SVZ	3.052	0.072	0.159	3.124
SUSY II	LCSR	4.148	0.266	0.460	4.413
	SVZ	3.054	0.084	0.167	3.138
SUSY III	LCSR	4.078	0.168	0.312	4.246
	SVZ	3.050	0.071	0.156	3.121

Table 8: Partial decay widths of $B \rightarrow K^* \tau^+ \tau^-$. Other conventions can be found at Table 7.

The partial decay widths (PDW) are listed in Tables. 5,6,7, and 8. We see that at the high $\hat{s}$ region, for the process $B \rightarrow K l^+ l^-$, $l=\mu, \tau$, the contributions of NHBs do show up, as expected. For $B \rightarrow K^* l^+ l^-$, the effects of NHBs in the high $\hat{s}$ region is significant when $l=\tau$ while they are small for $l=\mu$. It can be read out from these four tables that the results are consistent with the Fig. 1(a), 2(a), 3(a), and 4(a). In order to estimate the theoretical uncertainty brought by the methods calculating the weak form factors, we use the form factors calculated with LCSR and SVZ QCD sum rules (SVZ) method [17]. For $B \rightarrow K \ell^+ \ell^-$, PDWs calculated with form factors obtained by SVZ method is 50% of those by LCSR approach; while for $B \rightarrow K^* \ell^+ \ell^-$, PDWs increase 100% or so. We see that at low $\hat{s}$ regions the theoretical uncertainty can reach from 100% to 200%. Another point worthy of mention is that the contribution of resonances dominates the integrated decay width, as had been pointed out in [29].

5 Conclusion

We have calculated invariant mass spectrum, FBAs, and lepton polarizations for $B \rightarrow K \ell^+ \ell^-$ and $B \rightarrow K^* \ell^+ \ell^-$ $l=\mu, \tau$ in SUSY theories. In particular, we have analyzed the effects of NHBs to these processes. It is shown that the effects of the NHBs to $B \rightarrow K \tau^+ \tau^-$ and $B \rightarrow K^* \tau^+ \tau^-$ in some regions of parameter space of SUSY models are considerable and remarkable. The reason lies in the mass of the τ , which can magnify the effects of NHBs and can be seen through the related formula. The numerical results imply that there still exist possibilities to observe the effects of NHB in $B \rightarrow K \mu^+ \mu^-$ and $B \rightarrow K^* \mu^+ \mu^-$ through IMS, FBA and lepton polarizations of these processes. In particular, for $B \rightarrow K \mu^+ \mu^-$ in the case of SUSY II, the partial width in the high $\hat{s}$ where short distance physics dominates can be enhanced by a factor of 12 compared to SM. Our analysis also shows that the theoretical uncertainties brought in calculating of weak form factors are quite large. But the effects of NHBs will not be washed out and can stand out in some regions of the parameter space in MSSM. If only partial widths are measured, it is difficult to observe the effects of NHBs except for the decay $B \rightarrow K \tau^+ \tau^-$. However, the combined analysis of IBS, FBA, and lepton polarizations can provide useful knowledge to look for SUSY. Finally, we would like to point out that FBA for $B \rightarrow K l^+ l^-$ vanishes (or, more precisely, is negligibly small) in SM and it does not vanish in 2HDM and

SUSY models with large $\tan\beta$ due to the contributions of NHBs. However, only in SUSY models and for $l=\tau$ it is large enough to be observed in B factories in the near future.

Acknowledgements

One of the Authors (Q.S YAN) would like to thank Dr. W.J. HUO for his help during the work. This work was support in part by the National Nature Science Foundation of China.

References

- [1] B.Grinstein, M.J.Savage and M.B.Wise, *Nucl. Phys.* **B 319** (1989) 271.
- [2] T. Goto et.al., hep-ph/9812369; S. Baek and P. Ko, hep-ph/9904283; Y.G. Kim, P. Ko and J.S. Lee, *Nucl. Phys.* **B 544** (1999) 64, and references therein. For the earlier references, see, for example, the references in ref. [9].
- [3] Y.B. Dai, C.S. Huang and H.W. Huang, *Phys. Lett.* **B 390** (1997) 257.
- [4] C.S. Huang and Q.S. Yan, *Phys. Lett.* **B 442** (98) 209; C.S. Huang, W. Liao and Q.S. Yan, *Phys. Rev.* **D 59** (1999) 011701;
- [5] S. Fukae, C.S. Kim and T. Yoshikawa, hep-ph/9908229.
- [6] S. Fukae, C.S. Kim, T. Morozumi and T. Yoshikawa, *Phys. Rev.* **D 59** (1999) 074013
- [7] Y.G. Kim, P.Ko, and J.S. Lee, *Nucl. Phys.* **B 544** (1999) 64.
- [8] T. Goto *et.al.*, *Phys. Rev.* **D55** (1997) 4273; T. Goto, Y. Okada and Y. Shimizu, *Phys. Rev.* **D58** (1998) 094006.
- [9] E. Lunghi *et.al.*, Preprint hep-ph/9906286.
- [10] J. L. Hewett, *Phys. Rev.* **D 53** (1996) 4964.
- [11] Y. Grossman, Z. Ligeti and E. Nardi, *Phys. Rev.* **D 55** (1997) 2768.
- [12] K. Krüger and L.M. Sehgal, *Phys. Lett.* **B380** (1996) 199.
- [13] S. Choudhury et.al., hep-ph/9902355.
- [14] Chao-Shang Huang and Shou-Hua Zhu, *Phys. Rev.* **D 61** (2000) 015011.
- [15] I.I. Bigi, M. Shifman, N.G. Vrantsev and A.I. Vainstein, *Phys. Rev. Lett.* **71** (1993) 496; B. Blok, L. Kozrakh, M. Shifman and A.I. Vainstein, *Phys. Rev.* **D 49** (1994) 3356; A.V. Manohar and M.B. Wise, *Phys. Rev.* **D 49** (1994) 1310; S. Balk, T.G. Körner, D. Pirjol and K. Schilcher, *Z. Phys. C* 64 (1994) 37; A.F. Falk, Z. Ligeti, M. Neubert and Y. Nir, *Phys. Lett.* **B 326** (1994) 145.

- [16] W. Jaus and D. Wyler, *Phys. Rev.* **D 41** (1990) 3405; P. Colangelo et al., *Phys. Lett.* **B 317** (1993) 183; Ceng et al., *Phys. Rev.* **D 54** (1996) 3656.
- [17] P. Colangelo et al., *Phys. Rev.* **D 53** (1996) 3672; Erratum, *ibid*, **D 57** (1998) 3186.
- [18] I.I. Balitsky, V.M. Braun and A.V. Kolesnichenko, *Nucl. Phys.* **B312** (1989) 509.
- [19] V.L. Chernyak and I.R. Zhitnitsky, *Nucl. Phys.* **B345** (1990) 137.
- [20] V.M. Braun, Preprint NORDITA-98-1-P (hep-ph/9801222).
- [21] A. Khodjamirian and R. Rückl, Preprint WUE-ITP-97-049 (hep-ph/9801443).
- [22] P. Ball and V.M. Braun, *Phys. Rev.* **D58** (1998) 094016; *Nucl. Phys.* **B543** (1999) 201; P. Ball *et al.*, *Nucl. Phys.* **B529** (1998) 323; P. Ball, *JHEP* **09** (1998) 005; *JHEP* **01** (1999) 010.
- [23] A. Ali, P. Ball, L.T. Handoko, G. Hiller, hep-ph/9910221.
- [24] S. Glenn, *et.al.* (CLEO Collaboration), *Phys. Rev.* **D 60** (1999) 034017; T. Affolder, *et.al.* (CLEO Collaboration), hep-ex/9905004.
- [25] F. Krüger and J. C. Romão, hep-ph/0002089.
- [26] N.G. Deshpande and J. Trampetić, *Phys. Rev. Lett.* **60** (1988) 2583.
- [27] G. Burdman, hep-ph/9505352; W. Roberts, hep-ph/9512253.
- [28] D.S. Du and C. Liu, *Phys. Lett.* **B317** (1993) 179.
- [29] D.S. Liu hep-ph/9505314; M.R. Ahmady, D.S. Liu and A.H. Fariborz, hep-ph/9506235; M.R. Ahmady and D.S. Liu, hep-ph/9311335.
- [30] T.M. Aliev, C.S. Kim and Y.G. Kim, hep-ph/9910501.
- [31] M. Jezabek and P. Urban, hep-ph/9712440.
- [32] D. Melikhov, N. Nikitin, and S. Simula, hep-ph/9807464.
- [33] C. Greub, A. Ioannissian, and D. Wyler, *Phys. Lett.* **B 346** (1995) 149.
- [34] T.M. Aliev and M. Savci, hep-ph/0003188.
- [35] T. M. Aliev, M. Savci, A. Özpıneci, and H. Koru; *J. Phys. G* **24** (1998) 49
- [36] C.Q. Geng and C.P. Kao, *Phys. Rev.* **D54** (1996) 5636.
- [37] C.S. Huang *et al.*, hep-ph/9810412; *Commun. Theor. Phys.* **32** (1999) 499.
- [38] S. Bertolini, F. Borzynatu, A. Masiero and G. Ridolfi, *Nucl. Phys.* **B 353** (91) 591;

- [39] G. Buchalla, A.J. Buras, and M.E. Lauthenbacher, *Rev. Mod. Phys.* **68** (96) 1125; A. Buras, *et al.*, Nucl. Phys.B 4009337; *Nucl. Phys.* **B 400** (93) 75; S. Bertolini, *et al.*, *Nucl. Phys.* **B 353** (91) 591.
- [40] A. Ali, G.F. Giudice, and T. Mannel, *Z. Phys.* **C67** (1995) 417.
- [41] A.J. Buras and M. Münz, *Phys. Rev.* **D 52** (1995) 186.
- [42] A. Ali, T. Mannel and T. Morozumi, *Phys. Lett.* **B273** (1991) 505.
- [43] H.E.Logan and U.Nierste, hep-ph/0004139.